arXiv:math/0611431v3 [math.DG] 3 Dec 2006

INTEGRABILITY CRITERION FOR ABELIAN EXTENSIONS OF LIE GROUPS

PEDRAM HEKMATI

Abstract. We establish a criterion for when an abelian extension of infinite-dimensional Lie algebras $\hat{\mathfrak{g}} = \mathfrak{g} \oplus_\omega \mathfrak{a}$ integrates to a corresponding Lie group extension $A \hookrightarrow \widehat{G} \twoheadrightarrow G$, where G is connected, simply connected and $A \cong \mathfrak{a}/\Gamma$ for some discrete subgroup $\Gamma \subseteq \mathfrak{a}$. When $\pi_1(G) \neq 0$, the kernel A is replaced by a central extension $\widehat{A}$ of $\pi_1(G)$ by A.

1. Introduction

Given a group G with a normal subgroup N, one can construct the quotient group $H = G/N$. The theory of group extensions addresses the converse problem. Starting with H and N, what different groups G can arise containing N as a normal subgroup such that $H \cong G/N$? The problem can be formulated for infinite-dimensional Lie groups, but the situation is more delicate. Many familiar theorems break down and one must take into account topological obstructions. In particular, Lie's third theorem no longer holds and the question of integrability, i.e. whether a Lie algebra corresponds to a Lie group, becomes relevant.

The aim of this paper is to establish an integrability criterion for abelian extensions of infinite-dimensional Lie groups by generalizing a geometric construction for gauge groups, [5, 6, 7]. In sections 2 and 3 we review the basic definitions of infinite-dimensional Lie groups and their abelian extensions. Section 4 gives a detailed account of the construction which leads up to the integrability criterion.

2. Infinite-dimensional lie groups

We define infinite-dimensional Lie groups along the lines of [8], which should be consulted for further details and for concrete examples. The first step is to define the concept of an infinite-dimensional smooth manifold. Here the bottom line is to replace $\mathbb{R}^n$ (or $\mathbb{C}^n$) by

---

Date: November 25, 2006.

a more general model space on which a meaningful differential calculus can be developed. Essentially all familiar constructions in finite dimensions then carry over to the infinite-dimensional setting. We consider sequentially complete locally convex (s.c.l.c) topological vector spaces. These spaces have the property that every continuous path has a Riemann integral. We adopt the following notion of smoothness.

Definition. *Let E, F be s.c.l.c. topological vector spaces over $\mathbb{R}$ (or $\mathbb{C}$) and $f : U \to F$ a continuous map on an open subset $U \subseteq E$. Then f is said to be differentiable at $x \in U$ if the directional derivative*

$$df(x)(v) = \lim_{t\to 0} \frac{1}{t}(f(x+tv) - f(x))$$

exists for all $v \in E$. It is of class C^1 if it is differentiable at all points of U and

$$df : U \times E \to F,\ (x, v) \mapsto df(x)(v)$$

is a continuous map on $U \times E$. Inductively we say that f is of class C^n if df is a map of class C^{n-1} and of class C^∞ or smooth if it is of class C^n for all $n \geq 1$.

This definition coincides with the alternative notion of *convenient* smoothness [3] on Fréchet manifolds. A smooth manifold modeled on a s.c.l.c. topological vector space E is a Hausdorff topological space M with an atlas of local charts $\{(U_i, \phi_i)\}$ such that the transition functions are smooth on overlaps $U_i \cap U_j$. A Lie group G is a smooth manifold endowed with a group structure such that the operations of multiplication and inversion are smooth. The Lie algebra $\mathfrak{g}$ is defined as the space of left-invariant vector fields. A vector field $X : G \to TG$ is left-invariant if

$$(L_{g*}X)(h) = X(gh),\ \ \forall g, h \in G$$

where L_{g*} denotes the pushforward map of the diffeomorphism $L_g : G \to G,\ h \mapsto gh$. By definition X is completely determined by its value at the identity and $\mathfrak{g}$ is therefore identified with $T_{\mathbf{1}}G$ as topological vector spaces, endowed with the continuous Lie bracket of vector fields. The most striking feature of infinite-dimensional Lie theory is that results on existence and uniqueness of ordinary differential equations and the implicit function theorem do not hold beyond Banach Lie groups. Therefore a priori there is no exponential map and even if it exists, it does not have to be locally bijective. The existence and smoothness of the exponential function hinges on the notion of regularity.

Definition. *A Lie group G is called regular if for each $X \in C^\infty([0,1], \mathfrak{g})$, there exists $\gamma \in C^\infty([0,1], G)$ such that*

$$\gamma'(t) = L_{\gamma(t)*}(\mathbf{1}).X(t), \qquad \gamma(0) = \mathbf{1}$$

and the evolution map

$$\mathrm{evol}_G : C^\infty([0,1],\mathfrak{g}) \to G, \qquad X \mapsto \gamma(1)$$

is smooth.

In other words every smooth curve in the Lie algebra should arise, in a smooth way, as the left logarithmic derivative of a smooth curve in the Lie group. Note that regularity is a sharper condition than the requirement that the exponential map should be defined and smooth. Indeed if $\gamma(t)$ is the curve corresponding to the constant path $X(t) = X_0$ for some $X_0 \in \mathfrak{g}$, then $\gamma(1) = \exp(X_0)$. All known Lie groups modeled on s.c.l.c. topological vector spaces are regular [8]. In the convenient setting for calculus, it has been shown [4] that all connected regular abelian Lie groups are of the form $\mathfrak{a}/\Gamma$, for some discrete subgroup $\Gamma \subseteq \mathfrak{a}$ of an abelian Lie algebra $\mathfrak{a}$. Moreover, parallel transport exists for connections on principal bundles with regular structure group [2]. Important examples of regular Lie groups include gauge groups $C^\infty(M,G)$ and diffeomorphism groups $\mathrm{Diff}(M)$, where M is a smooth compact manifold and G is a finite-dimensional Lie group. From this point on we assume that the Lie groups are regular unless stated otherwise.
We digress to say a few words about the cohomology of Lie groups and Lie algebras.

2.1. **Lie group cohomology.** Let G be a Lie group. An abelian Lie group A is called a smooth G-module if there is a smooth G-action on A by automorphisms $G\times A \to A$, $(g,a) \mapsto g.a$. The set of smooth maps $f : G^n \to A$ such that $f(g_1,\dots,g_n) = 0$ whenever $g_j = \mathbf{1}$ for some j, are called *n-cochains* and form an abelian group $C^n(G,A)$ under pointwise addition. A cochain complex

$$\cdots \to C^{n-1}(G,A) \xrightarrow{\delta_{n-1}} C^n(G,A) \xrightarrow{\delta_n} C^{n+1}(G,A) \to \dots$$

is generated by the nilpotent homomorphisms $\delta_n : C^n(G,A) \to C^{n+1}(G,A)$ defined by

$$(\delta_n f)(g_1,\dots,g_{n+1}) =$$

$$= g_1.f(g_2,\dots,g_{n+1}) + \sum_{i=1}^{n}(-1)^i f(g_1,\dots,g_ig_{i+1},\dots,g_{n+1}) + (-1)^{n+1} f(g_1,\dots,g_n) \ .$$

Let $Z^n(G,A) = \ker\, \delta_n$ and $B^n(G,A) = \mathrm{im}\, \delta_{n-1}$ denote the subgroups of *n-cocycles* and *n-coboundaries* respectively. The n-th Lie cohomology group is given by

$$H^n(G,A) = \frac{Z^n(G,A)}{B^n(G,A)} \ .$$

2.2. **Lie algebra cohomology.** Let $\mathfrak{g}$ and $\mathfrak{a}$ be topological Lie algebras. Then $\mathfrak{a}$ is a continuous $\mathfrak{g}$-module if there is a continuous $\mathfrak{g}$-action, $\mathfrak{g} \times \mathfrak{a} \to \mathfrak{a}, (X, v) \mapsto X.v$. Denote by $C^n(\mathfrak{g}, \mathfrak{a})$ the vector space of continuous alternating multilinear maps $\omega : \mathfrak{g}^n \to \mathfrak{a}$. A cochain complex

$$\cdots \to C^{n-1}(\mathfrak{g}, \mathfrak{a}) \xrightarrow{d_{n-1}} C^n(\mathfrak{g}, \mathfrak{a}) \xrightarrow{d_n} C^{n+1}(\mathfrak{g}, \mathfrak{a}) \to \dots$$

is obtained by the nilpotent linear maps $d_n : C^n(\mathfrak{g}, \mathfrak{a}) \to C^{n+1}(\mathfrak{g}, \mathfrak{a})$ given by Palais' formula

$$(d_n\omega)(X_1, \dots, X_{n+1}) \equiv \sum_{i=1}^{n+1}(-1)^{i+1} X_i.\omega(X_1, \dots, \hat{X}_i, \dots, X_{n+1}) +$$

$$+ \sum_{i<j}(-1)^{i+j}\omega([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{n+1})$$

where $\hat{X}_i$ means that X_i is omitted. Let $Z^n(\mathfrak{g}, \mathfrak{a}) = \ker d_n$ and $B^n(\mathfrak{g}, \mathfrak{a}) = \operatorname{im} d_{n-1}$ denote the subspaces of *n-cocycles* and *n-coboundaries* respectively. The n-th Lie cohomology group is the quotient space

$$H^n(\mathfrak{g}, \mathfrak{a}) = \frac{Z^n(\mathfrak{g}, \mathfrak{a})}{B^n(\mathfrak{g}, \mathfrak{a})} .$$

In the context of Lie group and Lie algebra extensions, the second cohomology group H^2 is important in classifying topologically trivial abelian extensions as we will see. For $n \geq 2$ there is a derivation map $D_n : H^n(G, A) \to H^n(\mathfrak{g}, \mathfrak{a})$ given by [9]

$$(D_n f)(X_1, \dots, X_n) = \frac{\partial^n}{\partial t_1 \dots \partial t_n} \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) f(\gamma_{\sigma(1)}(t_{\sigma(1)}), \dots, \gamma_{\sigma(n)}(t_{\sigma(n)}))\Big|_{t_i=0}$$

where $\gamma_1(t_1), \dots, \gamma_n(t_n)$ is any set of smooth curves in G satisfying $\gamma_i(0) = \mathbf{1}$ and $\gamma_i'(0) = X_i \in \mathfrak{g}$.

3. Abelian extensions

Definition. *An extension of Lie groups is a short exact sequence with smooth homomorphisms*

$$\mathbf{1} \to A \xrightarrow{i} \widehat{G} \xrightarrow{p} G \to \mathbf{1}$$

such that p admits a smooth local section $s : U \to \widehat{G}$, $p \circ s = \mathrm{id}_U$, where $U \subset G$ is a local identity neighborhood.

The existence of a smooth local section means that $\widehat{G}$ is a principal A-bundle over G. The extension is called *abelian* if A is abelian and *central* if $i(A)$ lies in the center $Z(\widehat{G})$. Two extensions $\widehat{G}_1$ and $\widehat{G}_2$ are *equivalent* if there exists a smooth homomorphism $\phi : \widehat{G}_1 \to \widehat{G}_2$ which makes the following diagram commute:

$$\begin{tikzcd}
A \arrow[r, "i_1"] \arrow[d, "id_A"'] & \widehat{G}_1 \arrow[r, "p_1"] \arrow[d, "\phi"'] & G \arrow[d, "id_G"'] \\
A \arrow[r, "i_2"] & \widehat{G}_2 \arrow[r, "p_2"] & G
\end{tikzcd}$$

It is easy to verify by diagram chasing that ϕ must be an isomorphism. The definition for Lie algebras is analogue.

Definition. *An extension of topological Lie algebras is a short exact sequence with continuous homomorphisms*

$$\mathbf{0} \to \mathfrak{a} \xrightarrow{i} \hat{\mathfrak{g}} \xrightarrow{p} \mathfrak{g} \to \mathbf{0}$$

Two extensions $\hat{\mathfrak{g}}_1$ and $\hat{\mathfrak{g}}_2$ are said to be *equivalent* if there is an isomorphism of topological Lie algebras $\phi : \hat{\mathfrak{g}}_1 \to \hat{\mathfrak{g}}_2$ such that the following diagram commutes:

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "i_1"] \arrow[d, "id_{\mathfrak{a}}"'] & \hat{\mathfrak{g}}_1 \arrow[r, "p_1"] \arrow[d, "\phi"'] & \mathfrak{g} \arrow[d, "id_{\mathfrak{g}}"'] \\
\mathfrak{a} \arrow[r, "i_2"] & \hat{\mathfrak{g}}_2 \arrow[r, "p_2"] & \mathfrak{g}
\end{tikzcd}$$

Next we show how abelian extensions can be constructed explicitly. We have to make an assumption however. Viewed as a principal bundle, $\widehat{G}$ is assumed to be topologically trivial.

Proposition. *Let G be a Lie group, A a smooth G-module and $f \in Z^2(G, A)$ a 2-cocycle. The smooth manifold $G \times A$ endowed with the multiplication*

$$(g_1, a_1)(g_2, a_2) = (g_1 g_2, a_1 + g_1.a_2 + f(g_1, g_2))$$

defines an abelian extension $\widehat{G} = G \times_f A$ of G by A.

Associativity of the group law follows by the 2-cocycle property. The unit element is $(\mathbf{1}, 0)$ and $(g, a)^{-1} = (g^{-1}, -g^{-1}.(a + f(g, g^{-1})))$. The extension is topologically trivial since $s : G \to \widehat{G}, g \mapsto (g, f(g, g))$ defines a smooth global section. Moreover, the conjugation action of $\widehat{G}$ on A coincides with the smooth G-action. There is a similar cocycle construction for Lie algebras.

Proposition. *Let $\mathfrak{g}$ be a topological Lie algebra, $\mathfrak{a}$ a continuous $\mathfrak{g}$-module and $\omega \in Z^2(\mathfrak{g}, \mathfrak{a})$ a 2-cocycle. The topological vector space $\mathfrak{g} \oplus \mathfrak{a}$ endowed with the continuous Lie bracket*

$$[(X_1, v_1), (X_2, v_2)] = ([X_1, X_2], X_1.v_2 - X_2.v_1 + \omega(X_1, X_2))$$

defines a topologically split abelian extension $\hat{\mathfrak{g}} = \mathfrak{g} \oplus_\omega \mathfrak{a}$ *of* $\mathfrak{g}$ *by* $\mathfrak{a}$. *A continuous global section is given by* $s : \mathfrak{g} \to \hat{\mathfrak{g}}, X \mapsto (X, 0)$.

It turns out that all topologically trivial abelian extensions, where the conjugation action of $\widehat{G}$ on A induces the smooth G-action, arise in this way. Furthermore, two such extensions are equivalent if and only if the 2-cocycles differ by a 2-coboundary [9]. The second cohomology group H^2 therefore parametrize the set of equivalence classes of these extensions. The Lie algebra of $\widehat{G} = G \times_f A$ is as one would expect $\hat{\mathfrak{g}} = \mathfrak{g} \oplus_{D_2 f} \mathfrak{a}$.

4. INTEGRABILITY CRITERION

In this section we elucidate when an abelian extension of Lie algebras $\hat{\mathfrak{g}} = \mathfrak{g} \oplus_\omega \mathfrak{a}$ corresponds to a Lie group extension $\widehat{G}$. If $\omega = D_2 f$ for some $f \in Z^2(G, A)$, then by the previous section the corresponding Lie group extension is $\widehat{G} = G \times_f A$. In the general case, ω must satisfy a certain condition that will become apparent by the following construction. The basic idea is to construct $\widehat{G}$ as the quotient of a larger group $\mathcal{P}G \times_\gamma A$. This means that in general the extension will be topologically twisted and therefore the group multiplication cannot be described by a smooth global 2-cocycle.

Let G be the connected Lie group of $\mathfrak{g}$ and A a smooth G-module of the form $\mathfrak{a}/\Gamma$ for some discrete subgroup $\Gamma \subseteq \mathfrak{a}$. We write $e : \mathfrak{a} \to A$ for the exponential (quotient) map and employ multiplicative notation. The Lie algebra cocycle $\omega \in Z^2(\mathfrak{g}, \mathfrak{a})$ defines a closed G-equivariant 2-form $\omega^{eq} \in \Omega^2(G, \mathfrak{a})$ by

$$\omega^{eq}(g)(L_{g*}X, L_{g*}Y) = (L_g^* \omega^{eq})(\mathbf{1})(X, Y) = g.\omega(X, Y) \qquad \forall X, Y \in \mathfrak{g} \ .$$

For central extensions, the G-action on A is trivial and this is simply the associated left-invariant 2-form. Let $\mathcal{P}G$ denote the space of smooth based paths $\hat{g} : [0, 1] \to G$, originating at the identity $\hat{g}(0) = \mathbf{1}$ and ending at $\hat{g}(1) = g$. It is given the C^∞-topology of uniform convergence of the paths and all their derivatives. We require further that the left logarithmic derivative of the paths should be periodic, i.e. $\hat{g}^{-1}d\hat{g}(0) = \hat{g}^{-1}d\hat{g}(1)$. Consider $\mathcal{P}G \times A$ and introduce an equivalence relation

$$(\hat{g}_1, a_1) \sim (\hat{g}_2, a_1 e^{\int_{\pi[\hat{g}_1, \hat{g}_2]} \omega^{eq}})$$

whenever two paths end at the same point $\hat{g}_1(1) = \hat{g}_2(1)$ and form a smooth contractible loop, so that there is a well-defined 2-dimensional surface $\pi = \pi[\hat{g}_1, \hat{g}_2]$ in G bounded by these paths. If $\Delta_2 = \{(t, s) \in \mathbb{R}^2 | 0 \leq s \leq t \leq 1\}$, then one could take π as the smooth singular 2-chain

$$\pi : (t,s) \mapsto \begin{cases} g & t = 1,\ 0 \le s \le 1 \\ \hat{g}_1(\alpha(t(t-s)))\hat{g}_2(\alpha(ts)) & \text{otherwise} \end{cases}$$

where $\alpha : [0,1] \to [0,1]$ is any smooth map satisfying $\alpha(0) = 0, \alpha(1) = 1$ and $\frac{d\alpha}{du}(0) = \frac{d\alpha}{du}(1) = 0$. The surface π is not unique however. If π' is another smooth 2-chain with the same boundary, then

$$e^{\int_{\pi'} \omega^{eq}} = e^{\int_{\pi'} \omega^{eq} + \int_{\pi+\pi^-} \omega^{eq}} = e^{\int_{\pi'+\pi^-} \omega^{eq}} e^{\int_{\pi} \omega^{eq}}$$

where π^- denotes π with the opposite orientation. Here $\pi' + \pi^-$ is the smooth 2-cycle corresponding to the closed surface obtained by gluing together π' and π^- along their common boundary. For the equivalence relation to be well-defined we require that $e^{\int_{\pi'+\pi^-} \omega^{eq}} = 0$, or equivalently that

$$\int_c \omega^{eq} \in \Gamma \tag{1}$$

for all smooth 2-cycles $c \in Z_2(G)$. If $c = \partial b \in B_2(G)$ is a 2-boundary, then this is automatically satisfied by Stoke's theorem

$$\int_{\partial b} \omega^{eq} = \int_b d\omega^{eq} = 0$$

and therefore the condition factors through to homology.

We can now proceed to construct the Lie group extension $\hat{G}$ corresponding to $\hat{\mathfrak{g}} = \mathfrak{g} \oplus_\omega \mathfrak{a}$ by defining a multiplication on $\mathcal{P}G \times A/\sim$

$$[(\hat{g}_1, a_1)][(\hat{g}_2, a_2)] = [(\hat{g}_1\hat{g}_2, a_1(\hat{g}_1.a_2)e^{\gamma(\hat{g}_1,\hat{g}_2)})]$$

where $\hat{g}.a := \hat{g}(1).a = g.a$ is the given G-action on A and $\gamma : \mathcal{P}G \times \mathcal{P}G \to \mathfrak{a}$ is a smooth 2-cocycle. The latter must be defined in such a way that it yields the correct Lie algebra cocycle ω and is compatible with the equivalence relation. This is accomplished by choosing

$$\gamma(\hat{g}_1, \hat{g}_2) = \int_\sigma \omega^{eq}$$

where $\sigma : \Delta_2 \to G, (t,s) \mapsto \hat{g}_1(t)\hat{g}_2(s)$ is the smooth singular 2-chain with vertices in 1, g_1 and g_1g_2 and bounded by the paths $\hat{g}_1$, $\hat{g}_2$ and $g_1\hat{g}_2$. The 2-cocycle identity

$$(\delta_2\gamma)(\hat{g}_1, \hat{g}_2, \hat{g}_3) = \hat{g}_1.\gamma(\hat{g}_2, \hat{g}_3) - \gamma(\hat{g}_1\hat{g}_2, \hat{g}_3) + \gamma(\hat{g}_1, \hat{g}_2\hat{g}_3) - \gamma(\hat{g}_1, \hat{g}_2) = 0 \mod \Gamma$$

is satisfied by (1) since the regions of integration form a smooth 2-cycle, see Figure 1. The face not joining to $\mathbf{1}$ is the left translation by g_1 of the domain of integration

$$\hat{g}_1.\gamma(\hat{g}_2, \hat{g}_3) = \int_\sigma g_1.\omega^{eq} = \int_\sigma L^*_{g_1}\omega^{eq} = \int_{L_{g_1}\sigma} \omega^{eq}$$



Figure 1.

where we have used the G-equivariance of ω^{eq}. Thus, we conclude that the multiplication is associative. To see that it is well-defined, i.e. independent of the representatives, a straightforward calculation leads to

$$\int_{\pi[\hat{g}_1\hat{g}_2,\hat{g}_1'\hat{g}_2]}\omega^{eq}-\int_{\pi[\hat{g}_1,\hat{g}_1']}\omega^{eq}-\gamma(\hat{g}_1',\hat{g}_2)+\gamma(\hat{g}_1,\hat{g}_2)=0 \mod \Gamma$$

$$\int_{\pi[\hat{g}_1\hat{g}_2,\hat{g}_1\hat{g}_2']}\omega^{eq}-\int_{\pi[\hat{g}_2,\hat{g}_2']}g_1.\omega^{eq}-\gamma(\hat{g}_1,\hat{g}_2')+\gamma(\hat{g}_1,\hat{g}_2)=0 \mod \Gamma$$

$$\int_{\pi[\hat{g}_1\hat{g}_2,\hat{g}_1'\hat{g}_2']}\omega^{eq}-\int_{\pi[\hat{g}_1,\hat{g}_1']}\omega^{eq}-\int_{\pi[\hat{g}_2,\hat{g}_2']}g_1.\omega^{eq}-\gamma(\hat{g}_1',\hat{g}_2')+\gamma(\hat{g}_1,\hat{g}_2)=0 \mod \Gamma\ .$$

Again the regions of integration form closed 2-dimensional surfaces in G, depicted in Figure 2. The label on each face refers to the corresponding term in the expressions above, numbered from left to right.



Figure 2.

Next let us calculate the Lie algebra cocycle. We use the equivariance property to evaluate

ω^{eq} only at the identity

$$\begin{aligned}\int_\sigma \omega^{eq} &= \int_{\Delta_2} \omega^{eq}(\sigma(t,s))\left(\sigma_*\frac{\partial}{\partial t}, \sigma_*\frac{\partial}{\partial s}\right) dt\wedge ds \\ &= \int_{\Delta_2} \omega^{eq}(\hat{g}_1(t)\hat{g}_2(s))\left(\frac{d\hat{g}_1}{dt}\hat{g}_2(s), \hat{g}_1(t)\frac{d\hat{g}_2}{ds}\right) dtds \\ &= (g_1g_2).\int_{\Delta_2} \omega^{eq}(\mathbf{1})\left(\hat{g}_2^{-1}(s)\hat{g}_1^{-1}(t)\frac{d\hat{g}_1}{dt}\hat{g}_2(s), \hat{g}_2^{-1}(s)\frac{d\hat{g}_2}{ds}\right) dtds\ .\end{aligned}$$

Near the identity in $\mathcal{P}G$ we can write $\hat{g}_1(t) = \exp(\tau X(t))$ and $\hat{g}_2(s) = \exp(\sigma Y(s))$ for $X, Y \in \mathcal{P}\mathfrak{g}$, where exp is defined pointwise by the exponential map of G. The Lie algebra cocycle is given by

$$\begin{aligned}(D_2 e^\gamma)(X,Y) &= \frac{d^2}{d\sigma d\tau}(e^{\gamma(\hat{g}_1,\hat{g}_2)}e^{-\gamma(\hat{g}_2,\hat{g}_1)})|_{\tau=\sigma=0} \\ &= \int_{0\le s\le t\le 1} \omega^{eq}(\mathbf{1})\left(\frac{dX}{dt}, \frac{dY}{ds}\right) dsdt + \int_{0\le t\le s\le 1} \omega^{eq}(\mathbf{1})\left(\frac{dX}{dt}, \frac{dY}{ds}\right) dsdt \\ &= \int_{0\le s,t\le 1} \omega^{eq}(\mathbf{1})\left(\frac{dX}{dt}, \frac{dY}{ds}\right) dsdt = \omega^{eq}(\mathbf{1})(X(1),Y(1)) = \omega(X,Y)\end{aligned}$$

where we have used the antisymmetry of ω^{eq} and the fact that $\omega^{eq}(\mathbf{1})(X(0),Y(0)) = 0$. With that, we have verified that γ induces a well-defined group multiplication and the correct cocycle at the Lie algebra level. The Lie group extension corresponding to $\hat{\mathfrak{g}} = \mathfrak{g}\oplus_\omega \mathfrak{a}$ is the principal $\widehat{A}$-bundle $\widehat{G} = \mathcal{P}G \times_\gamma A/\sim\ \to G$ with the projection $[(\hat{g}, a)] \mapsto \hat{g}(1)$. The fiber $\widehat{A} = \pi_1(G)\times_\gamma A$ is a central extension of $\pi_1(G)$ by A. This is easily seen by the following. If $\pi_1(G) \hookrightarrow \tilde{G} \twoheadrightarrow G$ denotes the universal covering, then the same construction for $\tilde{G}$ gives

$$\begin{tikzcd} A \arrow[r, "i_1"] \arrow[d] & \mathcal{P}\tilde{G}\times_\gamma A/\sim \arrow[r, "p_1"] \arrow[d] & \tilde{G} \arrow[d, "q"'] \\ \widehat{A} \arrow[r, "i_2"] & \mathcal{P}G\times_\gamma A/\sim \arrow[r, "p_2"] & G \end{tikzcd}$$

Restriction to the subgroup $\pi_1(G)\subset\tilde{G}$ induces a central extension $A \xrightarrow{i_1} \pi_1(G)\times_\gamma A \xrightarrow{p_1} \pi_1(G)$, since $\pi_1(G)$ is discrete and acts trivially on A. Finally we have $\widehat{A} = \ker p_2 = \ker p_1\circ q = \pi_1(G)\times_\gamma A$. The right action of the structure group $\widehat{A}$ is $[(\hat{g},a)].[(\eta,a')] = [(\hat{g}\eta, a(\hat{g}.a')e^{\gamma(\hat{g},\eta)})]$.

To see that condition (1) is not only necessary but sufficient, let $A\hookrightarrow\widehat{G}\twoheadrightarrow G$ be a Lie group extension corresponding to $\hat{\mathfrak{g}} = \mathfrak{g}\oplus_\omega\mathfrak{a}$. We claim that ω^{eq} is the curvature form of a connection on this principal bundle. Indeed, if $pr : \mathfrak{g}\oplus_\omega\mathfrak{a}\to\mathfrak{a}$ denotes the projection onto the ideal $\mathfrak{a}$, then there is a canonical connection 1-form on $\widehat{G}$ given by

$$\alpha := -pr(g^{-1}dg) : \widehat{G}\to T\widehat{G}^*\otimes\mathfrak{a}$$

with the curvature $\Omega = d\alpha + \frac{1}{2}[\alpha, \alpha] = d\alpha$, where the last equality holds since $\mathfrak{a}$ is abelian. By the Maurer-Cartan equation

$$\Omega = -prd(g^{-1}dg) = \frac{1}{2}pr[g^{-1}dg, g^{-1}dg] \ .$$

We are interested in the pullback of the curvature 2-form onto the base space. Let $U \subset G$ be an identity neighborhood and define a smooth local section $s : U \to \widehat{G}$ by $g \mapsto (g, 0)$. Evaluating the pullback on $X, Y \in \mathfrak{g}$ at the identity, we get precisely

$$\begin{aligned} s^*(\Omega)(\mathbf{1})(X,Y) &= d\alpha((\mathbf{1},0))((X,0),(Y,0)) \\ &= \frac{1}{2}pr[(X,0),(Y,0)] \\ &= \frac{1}{2}\omega(X,Y) - \frac{1}{2}\omega(Y,X) = \omega(X,Y) \ . \end{aligned}$$

Equipped with a connection, one can define the horizontal lift of a curve on the base space and subsequently the notion of parallel transport. In particular, the holonomy around a loop $\eta : S^1 \to G$ is given by

$$\text{hol}(\alpha, \eta) = e^{\int_\eta \alpha} = e^{\int_\pi \omega^{eq}}$$

where π is a surface enclosed by η. The arbitrariness in the choice of this surface leads as before to the requirement $\int_{[c]} \omega^{eq} \in \Gamma$ for all smooth 2-cycles $[c] \in H_2(G)$. Thus, we are lead to the result:

Theorem (Integrability Criterion)**.** *Let G be a connected regular Lie group and A a smooth regular G-module of the form $\mathfrak{a}/\Gamma$ for some discrete subgroup $\Gamma \subseteq \mathfrak{a}$. The abelian Lie algebra extension $\hat{\mathfrak{g}} = \mathfrak{g} \oplus_\omega \mathfrak{a}$ integrates to a Lie group extension $\mathbf{1} \to \widehat{A} \to \widehat{G} \to G \to \mathbf{1}$ if and only if*

$$\int_{[c]} \omega^{eq} \in \Gamma$$

for all $[c] \in H_2(G)$, where $\widehat{A} = \pi_1(G) \times_\gamma A$ is a central extension of $\pi_1(G)$ by A.

When G is simply connected, then $\widehat{A} = A$ and by Hurewicz theorem $\pi_2(G) = H_2(G)$, so the condition coincides with that found in [9].

ACKNOWLEDGEMENTS

I would like to thank Jouko Mickelsson and Karl-Hermann Neeb for helpful comments.

REFERENCES

1. H. Glöckner, *Fundamental Problems in the Theory of Infinite-Dimensional Lie Groups.* J. Geom. Symm. Phys. **5** (2006), pp. 24-35.
2. A. Kriegl, P. W. Michor, *Regular Infinite Dimensional Lie Groups.* J. Lie Theory **7** (1997), pp. 61-99.
3. A. Kriegl, P. W. Michor, *The Convenient Setting of Global Analysis.* Amer. Math. Soc., Providence (1997).
4. P. W. Michor, J. Teichmann, *Description of infinite dimensional abelian regular Lie groups.* J. Lie Theory **9** (1999) pp. 487-489.
5. J. Mickelsson, *Current Algebras and Groups.* Plenum press, 233 Spring Street, New York (1989).
6. J. Mickelsson, R. Percacci, *Global Aspects of p-Branes.* J. Geom. Phys. **15** (1995), pp. 369-380.
7. M. K. Murray, *Another Construction of the Central Extension of the Loop Group.* Commun. Math. Phys. **116** (1988), pp. 73-80.
8. J. Milnor, *Remarks on Infinite-Dimensional Lie Groups.* "Relativité, Groupes et Topologie II" B. De-Witt and R. Stora (Eds), North-Holland, Amsterdam (1983), pp. 1007-1057.
9. K. H. Neeb, *Abelian extensions of infinite-dimensional Lie groups.* Mathematical works. Part XV. Luxembourg: Université du Luxembourg, Séminaire de Mathématique (2004), pp. 69-194.

MATHEMATICAL PHYSICS, ROYAL INSTITUTE OF TECHNOLOGY, SE-106 91 STOCKHOLM, SWEDEN

E-mail address: `pedram@kth.se`